

Evaluation for sustainable land use in mountain areas of Northwestern Yunnan Province, China

Jian Peng · Yanglin Wang · Jiansheng Wu ·
Qing Chang · Yuan Zhang

Received: 5 September 2006 / Accepted: 30 November 2006 / Published online: 8 February 2007
© Springer Science + Business Media B.V. 2007

Abstract As an important component of sustainable development in mountain areas, evaluation for sustainable land use is always one of the hotspots of researches on sustainable development. Traditional evaluation for sustainable land use mainly focuses on the sustainability of land use model and biological production on temporal scale, and overlooks the effects of land use patterns on the sustainability, while landscape ecology can be a good help to realize the spatial analysis of sustainable land use. In this study, a synthetic evaluation indexes system for sustainable land use was constructed through the application of landscape metrics. Taking Yongsheng County of Yunnan Province, China as a case study, a series of quantitative evaluation were conducted in 1996, 1999 and 2001, to monitor the temporal dynamics of regional land use sustainability. Two indicators, contributing amount of indexes, and obstacle amount of indexes, were also set up to ascertain the significance of all the evaluation indexes to the evaluation results. The results showed that, in the study phases, the land use sustainability of the whole county had been low with a stable but great spatial difference,

and great changes took place in regional land use system in 1999 with the deviation from the aim of sustainable land use. It also showed that, the most important indexes contributing for the land use sustainability in the study period, were the indexes of population density and land use degree, followed by the index of landscape diversity and cropping index. And the most important indexes counteracting the land use sustainability were the indexes of per unit area total production value of industry and agriculture, per unit area yield of cereal crops, landscape fragmentation, followed by the indexes of per unit area yield of economic crops and fertilizer consume per unit area.

Keywords Evaluation for sustainable land use · Contributing amount of indexes · Obstacle amount of indexes · Mountain areas · Northwestern Yunnan Province of China

Introduction

As an important basis of sustainable development, sustainable land use is always one of the key topics of researchers, policy-makers and the publics. With the definition of the criteria and standards of sustainable land use, evaluation for sustainable land use (ESLU) is the core of researches on sustainable land use. However, in the last decades, the researches on ESLU developed slowly, with an extensive basis on the five principles of sustainable land use proposed by FAO

J. Peng · J. Wu
Center of Digital City and Urban Landscape,
Shenzhen Graduate School, Peking University,
Shenzhen 518055, China

J. Peng · Y. Wang (✉) · J. Wu · Q. Chang · Y. Zhang
College of Environmental Sciences, Peking University,
Beijing 100871, China
e-mail: ylwang@urban.pku.edu.cn

(1993). It is in great need of other related disciplines to deepen the evaluation.

The sustainability of land use implies not only the sustainability of land use model and biological production on the temporal scale, but also includes the optimization of land use patterns on the spatial scale. However, traditional ESLU focusing on the social, economic or ecological benefits of regional land use, all can be categorized as the research on the temporal scale, lacking of the analysis of effects of spatial patterns on the land use sustainability (Peng et al. 2003). Taking spatial heterogeneity and ecological holism as its theoretical cores, landscape ecology can be a great help to a synthetic ESLU on the temporal and spatial scales, with a strong function in the analysis of the spatial patterns of regional land use (Peng et al. 2006; Wang and Yang 1999). However, although there were some researches that explored the combination between landscape ecology and sustainable land use or land management (Ericksen et al.

2002; Gulinck et al. 2001; Piorr 2003; Qiu and Fu 2000), or made a further discussion on the landscape sustainability or sustainable landscape (Antrop 2006; Botequilha and Ahern 2002; Paoletti 1999; Haines-Young 2000), few were conducted directly for ESLU in the view of landscape ecology, and only Peng et al. (2006) proposed a framework of landscape ecological evaluation for sustainable land use with the division of land use sustainability into such three aspects as landscape productivity, landscape threatening and landscape stability. But their study just proved the feasibility of the framework in measuring spatial differences of regional land use sustainability without testifying the ability in quantifying temporal dynamics of sustainable land use, and could not display the fundamental factors determining regional land use sustainability.

Therefore, with a case study in Yongsheng County of northwestern Yunnan Province, China, the objectives of the study reported in this paper are: (1) to understand the temporal dynamics and spatial differ-




Fig. 1 The study area in Yongsheng County, northwest Yunnan Province, China

ences of regional land use sustainability with the application of the framework of landscape productivity, landscape threatening, and landscape stability; (2) to ascertain the significance of all the evaluation indexes to the evaluation results through the introduction of two indicators, the contributing amount of indexes, and the obstacle amount of indexes; and (3) to make a conclusion on the evaluation indexes for sustainable land use with the comparison between this case study in mountain areas and that in coastal areas conducted by Peng et al. (2006).

Materials and methods

The study area

The study area, Yongsheng County in the northwest of Yunnan Province, China, locates between the longitude 100°22′–101°11′ east and between latitude 25°59′–27°04′ north, with Yangtze River flowing through the county (Fig. 1). The total area is 4,950 km² with 92.42% mountainous area. Yongsheng County has a low latitude and plateau monsoonal climate with clearly demarcated four seasons, drought in winter and spring while moisture in summer and autumn.

As the study area is located at the transition zone extending from the low altitude of Yunnan Plateau to the high altitude of Qinghai – Tibet Plateau, the natural conditions are quite poor with rugged terrain, lean soil, heavy water and soil erosion, low average annual precipitation about 1,000 mm, low average annual temperature between 7.9 and 10.5°C, and deficiency of light and heat resources. In a word, the

eco-environment of the study area is fragile.

The agricultural development in the study area has a certain foundations, but the second industry and the third industry have been stagnant for a few years. Until the end of 2001, the gross population of the whole county was 380,394, including 93% agricultural population and 30.49% minority population, which were major in Hui ethnic group, Yi ethnic group and etc. There are 6 towns and 12 villages in the whole county, that is, Yongbei, Jinguan, Liangguan, Qina, Renhe, and Chenghai towns, and Taoyuan, Pianjiao, Taiji, Shunzhou, Banqiao, Songping, Guanghua, Liude, Dongshan, Yangping, Daan and Dongfeng villages.

Evaluation indexes for sustainable land use

The sustainability of land use not only depends on the stabilization of land use patterns and the optimization of biological and non-biological productions from land use, but also is driven by human demands, which result in the pressure on regional land use. Generally speaking, the higher human demands for regional land use are, the higher the aim of sustainable land use is, and the lower the feasibility of sustainable land use is. Therefore, based on the method of Analytical Hierarchy Process (AHP), applying the theories of landscape ecology, the indexes system for evaluating regional sustainable land use can be constructed from three aspects: Landscape productivity, Landscape threatening and Landscape stability (Peng et al. 2006), as shown in Table 1. It’s hoped the indexes system will help to discover the distance between the aim of sustainable land use and the status quo of current land use, and the ability of achieving the

Table 1 The indexes for evaluating sustainable land use based on the method of AHP

Evaluation rule (weight)	Evaluation indexes (weight)
Landscape threatening (0.35)	Population density x_1 (0.125)
	Land use degree x_2 (0.125)
	Cropping index x_3 (0.100)
Landscape productivity (0.40)	Total production value of industry and agriculture per unit area x_4 (0.125)
	Yield of cereal crops per unit area x_5 (0.125)
	Yield of economic crops per unit area x_6 (0.075)
	Fertilizer consume per unit area x_7 (0.075)
Landscape stability (0.25)	Landscape diversity x_8 (0.100)
	Landscape fragmentation x_9 (0.075)
	Landscape contagion x_{10} (0.0375)
	Landscape fractal dimension x_{11} (0.0375)

sustainability aim in the temporal scale of human generation.

Landscape productivity reflects the capacity of land production, including biological productivity, economic benefits and potential yield of land use. The higher landscape productivity is, the more land production is, and the higher the possibility to realize sustainable land use is. As the cereal crops and economic crops are planted widely in mountain areas of Northwestern Yunnan Province, two indexes, yield of cereal crops per unit area and yield of economic crops per unit area, are chosen to measure the biological productivity of regional land use. Due to the poor natural conditions and infertile soil, the index of fertilizer consume per unit area is chosen to weigh the potential yield of regional land use. As a rule, the index of total production value of industry and agriculture per unit area is chosen to weigh the economic benefits of regional land use.

Landscape threatening is the pressure imposed on land use through human activities, which reflects human demands for land use. The more human demands from land use, the greater the pressure on land use is, the higher the aim of sustainable land use is, and the more difficulties there are to realize sustainable land use. Three indexes are chosen to evaluate landscape threatening, that is, population density, land use degree, and cropping index. The higher the value of the three indexes is, the higher the extent of landscape threatening is. Reflecting the degree of human land use, the index of land use degree is the area-weighted mean of grading index of different land use types in the whole area, and the grading index of unused land is 1, with forest land, grassland and water body 2, farmland and garden 3, and constructing land 4 (Liu 1992).

Landscape stability means the ability of keeping the stability of landscape patterns and functions. The higher landscape stability is, the stronger landscape resistance against external disturbance is, the stronger landscape resilience to ecological balance after disturbances is, and the more possibility there is to maintain spatial patterns and landscape functions. Generally speaking, in medium developed agricultural landscapes, the increase of landscape heterogeneity is good for the maintenance of landscape stability. According to landscape ecology, landscape patterns determine landscape functions. Four landscape metrics, that is, landscape diversity, landscape fragmentation, landscape contagion and landscape fractal dimension,

are chosen to measure the stability of landscape patterns. The bigger landscape diversity and landscape fractal dimension is, the stronger landscape stability is, while landscape fragmentation and landscape contagion are on the contrary. Definitions and more explanations on these metrics were given by Fu (1995), and Gustafson (1998).

According to AHP, four judgment matrixes with the level from 1 to 9, are constructed to calculate the weight of indexes, including the rule layer and index layer (Table 1). All the weights pass the consistency test.

Contributing amount of indexes and obstacle amount of indexes

The indexes system can realize the spatial comparison of land use sustainability between different spatial units and the temporal comparison between different periods of the same spatial unit. However, the indexes system cannot display the significance of indexes to the evaluation results. Therefore, two indicators, the contributing amount of indexes and the obstacle amount of indexes, are introduced (Peng et al. 2001). The meanings of the two indexes are as follows:

$$C_i = (A_i \times W_i) / \sum_{i=1}^n (A_i \times W_i) \times 100\%$$

$$P_i = (100 - A_i) \times W_i \sum_{i=1}^n [(100 - A_i) \times W_i] \times 100\%$$

where C_i is the contributing degree of index i to the evaluation results, P_i is the limiting degree of index i to the results, A_i is the standardization value of index i , W_i is the weight of index i , and n is the number of indexes. The determination of contributing factors and obstacle factors to the sustainability of regional land use may be realized through the ordering of C_i and P_i .

Source data and data processing

The data for this study are derived from two sources. On one hand, the values of six indexes, namely total production value of industry and agriculture per unit area, yield of cereal crops per unit area, yield of economic crops per unit area, fertilizer consume per unit area, population density and cropping index can be

found or calculated through *Statistics Yearbook of Yongsheng County in 1996, 1999 and 2001*. On the other hand, the values of the other five indexes, that is, land use degree, landscape diversity, landscape fragmentation, landscape contagion, and landscape fractal dimension are calculated through remote sensed images with the aid of landscape analysis software FRAGSTATS. Based on ERDAS 8.4 software, three LANDSAT-TM images (orbit 131/41 and 131/42, resolution 30 m) of February 1996, April 1999 and April 2001, are interpreted with the reference to field reconnaissance in July, 1999 and 2001, and land use map of Yongsheng County in 2000 (scale 1:75,000). As a result, six land use types are classified, that is, paddy field, dry land, forest land, shrubby grassland, water body and urban land.

The method of maximum difference normalization is introduced to carry out non-dimensional quantities of original data. Data standardization of all the indexes is as follows:

$$X'_{ij} = (X_{ij} - X_{j\min}) / (X_{j\max} - X_{j\min})$$

where X_{ij} is the original value of index j of evaluation unit i , X'_{ij} is the normalized value of X_{ij} , and $X_{j\max}$ and $X_{j\min}$ are the maximum and minimum of index j among all the evaluation units, respectively.

Results

Spatial difference of the sustainability of regional land use

Through the results of sustainable land use evaluation for all the 18 towns and villages of Yongsheng County in 1996, 1999 and 2001 as shown in Table 2, it could be concluded that, land use sustainability of the whole county in the study periods was relatively low, and behaved a trend of descending with a narrow range. Meanwhile, in the study period, the sustainability of land use of the county showed a high and stable spatial difference in the 18 towns and villages, as the maximum among the evaluation values of all the towns and villages was always about 1.5 times of the minimum.

Applying the statistic analysis software SPSS11.0 for windows, a hierarchical cluster analysis of evaluation values of all the 18 towns and villages in 1996, 1999, and 2001 was conducted. According to the cluster method of between-groups linkage, when 3.5 was taken as the squared Euclidean distance, all the 18 samples could be classified into five types (Fig. 2): (1) Towns I with the highest land use sustainability, including Chenghai town, Taiji village and Taoyuan village; (2) Towns II with higher land

Table 2 The evaluation results for sustainable land use in Yongsheng County in 1996, 1999 and 2001

Evaluation unit	Evaluation value in 1996	Evaluation value in 1999	Evaluation value in 2001
Yongsheng County	49.44	51.30	49.25
Yongbei Town	43.78	42.44	39.08
Renhe Town	51.58	52.24	53.62
Qina Town	54.51	51.96	52.13
Lianguan Town	49.85	48.87	46.82
Jinguan Town	50.00	47.50	38.49
Chenghai Town	63.30	65.19	60.96
Pianjiao Village	54.57	53.66	53.71
Taiji Village	58.83	59.89	57.54
Taoyuan Village	58.82	60.96	59.62
Shunzhou Village	43.80	44.99	46.14
Banqiao Village	47.26	44.37	46.45
Yangpin Village	44.48	49.25	43.28
Liude Village	44.88	43.48	40.23
Dongshan Village	40.50	41.90	43.36
Dongfeng Village	46.67	48.24	42.87
Daan Village	44.41	45.59	41.57
Songpin Village	42.13	44.50	41.53

use sustainability, including Renhe town, Qina town and Pianjiao village; (3) Towns III with moderate land use sustainability, including Liangguan town, Shunzhou village, Banqiao village, Yangpin village and Dongfeng village; (4) Towns IV with fluctuant and moderate land use sustainability, including Jinguan town and Guanghua village; (5) Towns V with lower land use sustainability, including Yongbei town, Liude village, Dongshan village, Daan village and Songpin village.

According to the difference of the dynamics of land use sustainability in the study period, all the 18 towns and villages could be classified into four types (Fig. 3): (1) Towns A with continual descending of land use sustainability, including Yongbei town, Liangguan town, Jinguan town and Liude village, which mainly resulted from the continual increasing of the values of landscape threatening indexes, such as the index of land use degree; (2) Towns B with continual ascending of land use sustainability, including Renhe town, Shunzhou village, Dongshan village and Guanghua village, which was mainly due to the continual increasing of the values of landscape



Fig. 2 Regional difference of land use sustainability in Yongsheng County during 1996–2001



Fig. 3 Regional difference of the change of land use sustainability in Yongsheng County during 1996–2001

productivity indexes, such as the indexes of yield of economic crops per unit area and yield of cereal crops per unit area; (3) Towns C with ascending following descending of land use sustainability, including Qina town, Pianjiao village and Banqiao village, which was mainly because of the change of the values of landscape productivity indexes, such as the index of yield of cereal crops per unit area; (4) Towns D with descending following ascending of land use sustainability, including Chenghai town, Taiji village, Taoyuan village, Yangpin village, Dongfeng village, Daan village and Songpin village, which mainly resulted from the change of both landscape productivity indexes and landscape stability indexes, such as the indexes of yield of economic crops per unit area and landscape fractal dimension.

It was remarkable that, from 1996 to 2001, there were 13 towns and villages occupying 69.13% area and 76.96% population of the whole county, that had a decline of land use sustainability; while from 1996 to 1999, there were 11 towns and villages occupying 67.77% area and 47.45% population of the whole county, that had an increase of land use sustainability.

Therefore, it could be concluded that, regional land use system changed greatly in 1999, which deviated from the aim of sustainable land use.

Contributing indexes and obstacle indexes of sustainable land use

Setting no less than 10% as the standard, according to the contributing amount and obstacle amount of the 11 evaluation indexes, the dominant contributing indexes and obstacle indexes of all the 18 towns and villages were ascertained, respectively. As shown in Table 3, it could be found that: (1) The dominant contributing indexes changed little from 1996 to 2001, because there were 13 towns and villages without the change of the primary contributing index, and the first two contributing indexes didn't change in all the towns and villages; (2) The dominant obstacle indexes also changed little from 1996 to 2000, as there were 12 towns and villages without the change of the primary obstacle index, and the first two obstacle indexes didn't change in 11 towns and villages; (3) It was the index of population density

and land use degree that were the most important contributing indexes to sustainable land use in the study period, followed by the index of landscape diversity and cropping index; and (4) Total production value of industry and agriculture per unit area, yield of cereal crops per unit area and landscape fragmentation were the most important obstacle indexes, followed by the indexes of yield of economic crops per unit area and fertilizer consume per unit area.

Through the analysis of the contributing amount and obstacle amount of evaluation indexes for sustainable land use, it could be concluded that, although land use systems of the 18 towns and villages in Yongsheng County have changed in different ways during the study periods, the contributing amount and obstacle amount of evaluation indexes to the evaluation results of land use sustainability in each town or village, were relatively stable. Actions could be taken on the selected dominant contributing indexes and obstacle indexes, so as to optimize spatial patterns and to perfect landscape functions of regional land use systems. Generally speaking, there were mainly two ways to enhance regional land use sustainability. One was to increase

Table 3 The dominant contributing indexes and obstacle indexes for evaluating sustainable land use in Yongsheng County

Towns (T)/ Villages (V)	The dominant contributing indexes			The dominant obstacle indexes		
	1996	1999	2001	1996	1999	2001
Yongbei T	x ₈ , x ₂ , x ₄ , x ₅ , x ₁	x ₈ , x ₄ , x ₂ , x ₅ , x ₁	x ₈ , x ₄ , x ₅ , x ₁	x ₁ , x ₃ , x ₅ , x ₉ , x ₆ , x ₄ , x ₇	x ₃ , x ₁ , x ₅ , x ₂ , x ₆ , x ₉ , x ₄ , x ₇	x ₂ , x ₁ , x ₃ , x ₉ , x ₅ , x ₄
Renhe T	x ₁ , x ₂ , x ₅ , x ₈	x ₁ , x ₂ , x ₅	x ₁ , x ₂ , x ₅ , x ₈	x ₄ , x ₉ , x ₃	x ₄ , x ₉ , x ₃ , x ₈	x ₄ , x ₃ , x ₉
Qina T	x ₅ , x ₈ , x ₁ , x ₂	x ₅ , x ₈ , x ₁	x ₅ , x ₈ , x ₁ , x ₇ , x ₂	x ₄ , x ₉ , x ₂ , x ₃ , x ₁	x ₄ , x ₃ , x ₂ , x ₉ , x ₁	x ₃ , x ₄ , x ₂ , x ₉ , x ₁
Liangguan T	x ₅ , x ₄ , x ₈ , x ₂	x ₄ , x ₅ , x ₈	x ₄ , x ₅ , x ₈	x ₁ , x ₂ , x ₃ , x ₉	x ₁ , x ₂ , x ₃ , x ₉ , x ₆	x ₁ , x ₂ , x ₃ , x ₉ , x ₆
Jinguan T	x ₅ , x ₄ , x ₈ , x ₂	x ₅ , x ₄ , x ₈	x ₅ , x ₈ , x ₄	x ₁ , x ₃ , x ₂ , x ₇	x ₁ , x ₂ , x ₃ , x ₇	x ₂ , x ₁ , x ₃ , x ₆ , x ₉
Chenghai V	x ₅ , x ₂ , x ₁ , x ₈ , x ₃	x ₅ , x ₁ , x ₈ , x ₂ , x ₃	x ₅ , x ₈ , x ₁ , x ₂ , x ₃	x ₄ , x ₉ , x ₇	x ₄ , x ₉	x ₄ , x ₉ , x ₆
Pianjiao V	x ₁ , x ₂ , x ₅ , x ₈ , x ₃	x ₁ , x ₈ , x ₅ , x ₃ , x ₂	x ₁ , x ₈ , x ₅ , x ₃ , x ₂	x ₄ , x ₉ , x ₇	x ₄ , x ₉ , x ₂	x ₄ , x ₂ , x ₉
Taiji V	x ₂ , x ₁ , x ₅ , x ₆	x ₁ , x ₂ , x ₅ , x ₆ , x ₈	x ₁ , x ₂ , x ₅ , x ₃	x ₄ , x ₉ , x ₈ , x ₇ , x ₃	x ₄ , x ₉ , x ₃	x ₄ , x ₉ , x ₅ , x ₈
Taoyuan V	x ₁ , x ₅ , x ₃ , x ₂ , x ₈	x ₅ , x ₁ , x ₃ , x ₈ , x ₆	x ₅ , x ₁ , x ₃ , x ₈ , x ₇ , x ₆	x ₄ , x ₉ , x ₂	x ₄ , x ₂ , x ₉ , x ₇	x ₄ , x ₂ , x ₉
Shenzhou V	x ₁ , x ₃ , x ₈ , x ₂	x ₁ , x ₃ , x ₂ , x ₈	x ₁ , x ₂ , x ₃ , x ₈	x ₄ , x ₅ , x ₆ , x ₉ , x ₇ , x ₂	x ₅ , x ₄ , x ₉ , x ₇ , x ₆	x ₄ , x ₅ , x ₉ , x ₆ , x ₇
Banqiao V	x ₁ , x ₂ , x ₃ , x ₅ , x ₈	x ₁ , x ₂ , x ₈ , x ₃	x ₁ , x ₂ , x ₈	x ₄ , x ₅ , x ₆ , x ₉ , x ₇	x ₄ , x ₅ , x ₆ , x ₉ , x ₇	x ₄ , x ₅ , x ₆ , x ₃
Yangpin V	x ₁ , x ₂ , x ₃ , x ₈	x ₁ , x ₂ , x ₃ , x ₈	x ₂ , x ₁ , x ₃	x ₅ , x ₄ , x ₆ , x ₇ , x ₉	x ₄ , x ₅ , x ₆ , x ₇ , x ₉	x ₅ , x ₄ , x ₇ , x ₆ , x ₈
Liude V	x ₁ , x ₂ , x ₃ , x ₈	x ₁ , x ₂ , x ₃ , x ₈	x ₁ , x ₂ , x ₈ , x ₃	x ₄ , x ₅ , x ₇ , x ₉	x ₄ , x ₅ , x ₉ , x ₇	x ₄ , x ₅ , x ₆ , x ₉
Dongshan V	x ₁ , x ₂ , x ₉	x ₁ , x ₂ , x ₉ , x ₃	x ₁ , x ₂ , x ₃ , x ₉	x ₅ , x ₄ , x ₈ , x ₇ , x ₃	x ₄ , x ₅ , x ₈ , x ₇ , x ₆	x ₄ , x ₅ , x ₈ , x ₇ , x ₆
Dongfeng V	x ₁ , x ₂ , x ₃ , x ₈	x ₁ , x ₂ , x ₃	x ₁ , x ₂ , x ₃ , x ₈	x ₄ , x ₅ , x ₇	x ₄ , x ₅ , x ₇ , x ₈	x ₄ , x ₅ , x ₇
Daan V	x ₁ , x ₂ , x ₈ , x ₃	x ₁ , x ₂ , x ₈ , x ₃	x ₁ , x ₂ , x ₈ , x ₃	x ₄ , x ₅ , x ₇ , x ₆	x ₄ , x ₅ , x ₉ , x ₇	x ₄ , x ₅ , x ₉ , x ₇
Songpin V	x ₂ , x ₁	x ₁ , x ₂ , x ₈	x ₁ , x ₂ , x ₈	x ₄ , x ₅ , x ₇ , x ₃ , x ₈	x ₄ , x ₅ , x ₇ , x ₃ , x ₆	x ₄ , x ₅ , x ₃ , x ₇
Guanghua V	x ₁ , x ₂ , x ₈	x ₁ , x ₂ , x ₈ , x ₅	x ₁ , x ₈ , x ₆ , x ₂ , x ₅	x ₄ , x ₅ , x ₃ , x ₇ , x ₆ , x ₉	x ₄ , x ₃ , x ₆ , x ₉ , x ₅ , x ₇	x ₄ , x ₃ , x ₅ , x ₂ , x ₉

Indexes with the contributing amount or obstacle amount of no less than 10% are listed. The order of indexes in the table indicates the magnitude of the contributing or obstacle amount of indexes, and indexes with higher contributing or obstacle amount rank first.

the contributing amount of indexes, such as limiting population growth, controlling the construction in the towns and villages, regulating cropping index, and enhancing landscape diversity. And the other was to decrease the obstacle amount of indexes, such as increasing landscape productivity, and decreasing landscape fragmentation.

Discussion

Selection of evaluation indexes

The evaluation indexes system for regional sustainable land use is the core of researches on the evaluation for sustainable land use (Chen 2001; Chen and Zhang 2001). As there are too many indexes that can indicate the sustainability of land use, and the difference of the aim of sustainable land use in different areas makes different evaluation emphases, regional dominance must be regarded as the most important principle in selecting evaluation indexes for sustainable land use. For example, in this case study, the indexes of yield of cereal crops per unit area and yield of economic crops per unit area are selected to indicate landscape productivity, while in the case study in coastal areas conducted by Peng et al. (2006), the indexes of yield of cereal crops per unit area and yield of aquaculture per unit area are chosen. It is mainly owing to regional differences. In mountain areas of Northwestern Yunnan Province, China, agricultural land is mainly utilized for the planting of cereal and economic crops, while in coastal areas of Northwestern Shandong Province, China, cereal crops planting and aquaculture are the main agricultural activities.

Efficacy of evaluation indexes

The index of landscape fragmentation is chosen to evaluate landscape stability both in this case study and that in coastal areas conducted by Peng et al. (2006), but the efficacies of the index are on the contrary, one positive and the other negative to land use sustainability. It may be due to regional difference of landscape matrix. In mountain areas of northwestern Yunnan Province, China, the deep ravines lead to high fragmented topography. The high heterogeneity of mountain landscape in Yongsheng County, contrasts clearly with the low heterogeneity of coastal land-

scape in northwestern Shandong Province, China. According to the theory of moderate disturbances in landscape ecology, the increase of landscape fragmentation in low heterogeneous landscapes will help to the increase of landscape heterogeneity and the enhancement of landscape stability. Reversely, the increase of landscape fragmentation in high heterogeneous landscapes will block the flows of function, material and information in the landscape, and go against the maintenance of the stability of landscape spatial patterns and functions.

Weight of evaluation indexes

The method of AHP is often used to quantify the weights of evaluation indexes for sustainable land use. Recently, some other objective methods such as regression coefficient method and mean square method are used (Liu and Hao 2003; Zhang et al. 2002). But in these methods, the weights of indexes are all inevitably influenced by the actual values of indexes, and it is difficult to really reflect the relative difference of the importance of indexes. Therefore, aiming at reflecting the difference of relative importance of indexes, the method of AHP is prior to other methods.

Threshold of evaluation indexes

The threshold of indexes is an important parameter of judging if the index is sustainable or how much the sustainability is. Generally speaking, scientific and reasonable threshold should be obtained through control experiments, and should not refer to experiential planning value or target value, which cannot correlate to the sustainability of land use directly. As a result, if we cannot determinately quantify the target of sustainable land use, the aim of sustainable land use evaluation should be set as the discussion on the temporal dynamics or spatial differences of land use sustainability. In this sense, it is useless to discuss whether a single regional land use system is sustainable or not.

Conclusions

Traditional ESLU with the social, economic or ecological benefits of regional land use can be

categorized as the research on the temporal scale, lacking of the evaluation for spatial patterns of sustainable land use (Peng et al. 2003). It is of great significance to incorporate land use pattern analysis into the evaluation for sustainable land use, according to the concepts and methods of landscape ecology. The study reported in this paper proved the feasibility of the framework of landscape productivity, landscape threatening and landscape stability in evaluating sustainable land use in mountain areas. The results also showed that, land use sustainability of the study area in the study periods had been low with a high and stable spatial difference, and regional land use systems changed greatly in 1999 which deviated from the aim of sustainable land use. In the study periods, the indexes of population density and land use degree, followed by landscape diversity and cropping index orderly, were the dominant contributing indexes to sustainable land use. The indexes of total production value of industry and agriculture per unit area, yield of cereal crops per unit area, and landscape fragmentation, followed by yield of economic crops per unit area and fertilizer consume per unit area, were the dominant obstacle indexes to sustainable land use.

Acknowledgements The research is supported by National Natural Science Foundation of China (40635028, 40571051, 40471002). The authors wish to thank the anonymous reviewers for their reviews and helpful comments.

References

- Antrop, M. (2006). Sustainable landscapes: Contradiction, fiction or utopia? *Landscape and Urban Planning*, 75, 187–197.
- Botequilha, L. A., & Ahern, J. (2002). Applying landscape ecological concepts and metrics in sustainable landscape planning. *Landscape and Urban Planning*, 59, 65–93.
- Chen, B. M. (2001). A blue print of land use region based on regional indicator system for sustainable land use in China. *Progress in Geography*, 20, 247–253 (in Chinese).
- Chen, B. M., & Zhang, F. R. (2001). Theory and methodology for sustainable land use indicator system in China. *Journal of Natural Resources*, 16, 197–203 (in Chinese).
- Erickson, P. J., McSweeney, K., & Madison, F. W. (2002). Assessing linkages and sustainable land management for hillside agroecosystems in Central Honduras: Analysis of intermediate and catchment scale indicators. *Agriculture, Ecosystems and Environment*, 91, 295–311.
- FAO. (1993). FESLM: *An international framework for evaluating sustainable land management*. World Soil Resources Report, 73.
- Fu, B. J. (1995). The spatial pattern analysis of agricultural landscape in the loess area. *Acta Ecologica Sinica*, 15, 113–120 (in Chinese).
- Gulinck, H., Mugica, M., de Lucio, J. V., & Atauri, J. A. (2001). A framework for comparative landscape analysis and evaluation based on land cover data, with an application in the Madrid region (Spain). *Landscape and Urban Planning*, 55, 257–270.
- Gustafson, E. J. (1998). Quantifying landscape spatial pattern: what is the state of the art? *Ecosystem*, 1, 143–156.
- Haines-Young, R. (2000). Sustainable development and sustainable landscapes: Defining a new paradigm for landscape ecology. *Fennia*, 178, 7–14.
- Liu, J. Y. (1992). *The utilization of land in Tibet*. Beijing: Science Press (in Chinese).
- Liu, Y. Q., & Hao, J. M. (2003). Evaluation on spatial differentia of regional sustainable land use. *Resources Science*, 25, 56–62 (in Chinese).
- Paoletti, M. G. (1999). Using bioindicators based on biodiversity to assess landscape sustainability. *Agriculture, Ecosystem, and Environment*, 74, 1–18.
- Peng, B. Z., An, X. D., Chen, F., & Pu, L. J. (2001). A study on sustainable land use in the Yangtze River Delta. *Journal of Natural Resources*, 16, 305–312 (in Chinese).
- Peng, J., Wang, Y. L., Li, W. F., Yue, J., Wu, J., & Zhang, Y. (2006). Evaluation for sustainable land use in coastal areas: A landscape ecological prospect. *International Journal of Sustainable Development and World Ecology*, 13, 25–36.
- Peng, J., Wang, Y. L., Song, Z. Q., Jing, J., & Ding, Y. (2003). Research progress on evaluation for sustainable land use. *Resources Science*, 25, 85–93 (in Chinese).
- Pierr, H. P. (2003). Environmental policy, agri-environmental indicators and landscape indicators. *Agriculture, Ecosystems and Environment*, 98, 17–33.
- Qiu, Y., & Fu, B. J. (2000). Land evaluation for sustainable use based on landscape ecologic theory. *Resources Science*, 22, 1–8 (in Chinese).
- Wang, Y. L., & Yang, X. J. (1999). The research on sustainable tourism development in the overall planning of scenic resort. *Resources Science*, 21, 37–43 (in Chinese).
- Zhang, F. R., Qi, W., Xue, Y. S., & Xu, L. (2002). The methods for evaluating sustainable land management. *Journal of China Agricultural University*, 7, 40–46 (in Chinese).